

ORAL PRESENTATION

Open Access

# Learning from the dead: improving safety while placing unconscious trauma patients in various lateral positions

Per Kristian Hyldmo<sup>1,2\*</sup>, Bryan P Conrad<sup>3</sup>, Dewayne N Dubose<sup>3</sup>, Jo Røislien<sup>1,6</sup>, Mark Prasarn<sup>4</sup>, Eldar Søreide<sup>2,7</sup>, Glenn Rechten<sup>5</sup>, MaryBeth Horodyski<sup>3</sup>

From London Trauma Conference 2013  
London, UK. 10-13 December 2013

## Background

The unconscious trauma patient with a possible unstable spinal injury constitutes a clinical challenge. To protect the unintubated airway, some guidelines [1,2] recommend that the patient be turned into a lateral position, e.g. the Recovery Position (RP) [1] or the Lateral Trauma Position (LTP) [2]. Other lateral positions have also been proposed, as the HAINES position [3] and variations thereof. However, moving the patient may cause secondary neurological injury. The aim of this study was to explore how much motion lateral position techniques produce in an unstable cervical spine injury.

## Method

We surgically created a global ligamentous instability between C5 and C6 in five fresh cadavers [4]. Four different techniques were evaluated; RP, LTP and two varieties of HAINES (one or both legs flexed; H1 and H2). Relative angular and linear motion between C5 and C6 was measured using an electromagnetic tracking device (Liberty, Polhemus Inc.™, Colchester, VT). Each method was repeated three times in each cadaver. Both angular and linear movements were measured. Data were analysed using generalized linear mixed models (GLMM), adjusting for intra-cadaver correlation.

## Results

Compared to RC, LTP created significantly less movement during lateral bending ( $p=.037$ ), while H1 and H2 had significantly less movement than RC in axial translation ( $p=.009$  and  $.033$ ). There was a tendency towards

LTP and H1 and H2 performing better than RC also for other movements.

## Conclusion

Our results indicate that in unconscious trauma patients, LTP or one of the two HAINES techniques is preferable to the classic recovery position in the setting of an unstable cervical spine injury.

## Authors' details

<sup>1</sup>Research Department, Norwegian Air Ambulance Foundation, Drøbak, Norway. <sup>2</sup>Network for Medical Sciences, University of Stavanger, Stavanger, Norway. <sup>3</sup>Department of Orthopaedics & Rehabilitation, University of Florida, Gainesville, Florida, USA. <sup>4</sup>Department of Orthopaedics, University of Texas, Huston, Texas, USA. <sup>5</sup>Associate Chief of Staff, Bay Pines VAHCS, St. Petersburg, Florida, USA. <sup>6</sup>Department of Biostatistics, University of Oslo, Oslo, Norway. <sup>7</sup>Department of Anaesthesiology and Intensive Care, Stavanger University Hospital, Stavanger, Norway.

Published: 7 July 2014

## References

1. Deakin CD, et al: *European Resuscitation Council Guidelines for Resuscitation 2010 Section 4. Adult advanced life support. Resuscitation* 2010, **81**:1305-1352.
2. Berlac P, et al: *Pre-hospital airway management: guidelines from a task force from the Scandinavian Society for Anaesthesiology and Intensive Care Medicine. Acta Anaesthesiol Scand* 2008, **52**:897-907.
3. Haines J: *Positioning an unconscious patient with suspected neck injury. JEMS: Journal of Emergency Medical Services* 1996, **21**:85-85.
4. Horodyski M, et al: *Cervical Collars are Insufficient for Immobilizing an Unstable Cervical Spine Injury. J Emerg Med* 2011, **41**(5):513-519.

doi:10.1186/1757-7241-22-S1-O4

**Cite this article as:** Hyldmo et al.: Learning from the dead: improving safety while placing unconscious trauma patients in various lateral positions. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine* 2014 **22**(Suppl 1):O4.

<sup>1</sup>Research Department, Norwegian Air Ambulance Foundation, Drøbak, Norway

Full list of author information is available at the end of the article