


CORRECTION

Open Access



Correction to: Symptoms presented during emergency telephone calls for patients with spontaneous subarachnoid haemorrhage

Asger Sonne^{1*} , Sarita Egholm¹, Laurits Elgaard¹, Niklas Breindahl¹, Alice Herrlin Jensen¹, Vagn Eskesen^{2,7}, Freddy Lippert^{3,7}, Frans Boch Waldorf^{4,5}, Nicolai Lohse^{6,7} and Lars Simon Rasmussen^{1,7}

Correction to: Scand J Trauma Resusc Emerg Med (2021) 29:118

<https://doi.org/10.1186/s13049-021-00934-x>

Following the publication of the original article [1], the authors informed as of an error in Figure 2: The term “persistently unconscious” had accidentally replaced

“neck pain” in the third row under “Symptoms”.

The correct figure is shown here below and has already been included in the original article.

The original article can be found online at <https://doi.org/10.1186/s13049-021-00934-x>.


*Correspondence: Asger.sonne@regionh.dk

¹ Department of Anaesthesia, Section 6011, Center of Head and Orthopaedics, Rigshospitalet, Inge Lehmanns Vej 6, 2100 Copenhagen, Denmark

Full list of author information is available at the end of the article



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



Author details

¹Department of Anaesthesia, Section 6011, Center of Head and Orthopaedics, Rigshospitalet, Inge Lehmanns Vej 6, 2100 Copenhagen, Denmark. ²Department of Neurosurgery, The Neuroscience Centre, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark. ³Copenhagen Emergency Medical Services, Copenhagen, Denmark. ⁴Research Unit of General Practice, Department of Public Health, University of Southern Denmark, Odense, Denmark. ⁵The Research Unit for General Practice and Section of General Practice, Department of Public Health, University of Copenhagen, Copenhagen, Denmark. ⁶Department of Emergency Medicine, Copenhagen University Hospital – Nordsjællands Hospital, Hillerød, Denmark. ⁷Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark.

Published online: 14 October 2021

Reference

1. Sonne, et al. Symptoms presented during emergency telephone calls for patients with spontaneous subarachnoid haemorrhage. *Scand J Trauma Resusc Emerg Med*. 2021;29:118. <https://doi.org/10.1186/s13049-021-00934-x>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

