

Oral presentation

Open Access

Interhospital transport of critically ill patient with dedicated ICU ventilator

R Vacchi, E Santoro, A Giugni* and P Cavallo

Address: Department of Emergency-Urgency Medicine, Intensive Care Unit, Maggiore Hospital, Bologna, Italy

Email: A Giugni* - aimonegiugni@gmail.com

* Corresponding author

from Scandinavian Update on Trauma, Resuscitation and Emergency Medicine 2009 Stavanger, Norway. 23 – 25 April 2009

Published: 28 August 2009

Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine 2009, 17(Suppl 3):O21 doi:10.1186/1757-7241-17-S3-O21

This abstract is available from: <http://www.sjtreem.com/content/17/S3/O21>

© 2009 Vacchi et al; licensee BioMed Central Ltd.

Introduction

Emilia Romagna is a region characterized by the presence of Three Integrated Trauma Systems Care (ITSC). Severe, resource-consuming patients are centralized in first level Hubs to guarantee high quality and efficient care. Every ITSC has a Trauma Center (Hub) connected with a net of peripheral hospitals (Spokes). A Helicopter Emergency Medical Service (HEMS) has been active since June 14 1986 at the Ospedale Maggiore, it covers most of the Bologna and Modena region and part of Ferrara, in addition to this, a new Trauma Center specialized in pediatric patients is developing outside the region. This means transfer times of at least 50 to 60 minutes by helicopter to consent patients to move from an Intensive Care Unit (ICU) to another. Obviously the HEMS play a fundamental role in such a setting.

Methods

Our HEMS counts numerous interhospital transfers of adult patients with respiratory failure and pediatric patients that needed respiratory assistance with high performance ventilators. Patients with severe respiratory failure may have to remain on an ICU ventilator throughout the whole transport period anyway transport ventilators may not be adequate for certain age groups in pediatric patients. We would like to show how we faced the necessity of transporting acute but stable patients, using a system that enables us to embark a high performance ventilator and a multi-parametric monitor in a safe and quick manner. We will discuss the equipment assemblage, methods

and techniques of intensive care in the critical patient in this setting.

Conclusion

Our experienced medical team has demonstrated to safely transport even the most critically ill patients if the care is optimized before departure.

References

1. Beninati W, Meyer MT, Carter TE: **The critical care air transport program.** *Crit Care Med* 2008, **36(7 Suppl)**:S370-376.
2. Kashani KB, Farmer JC: **The support of severe respiratory failure beyond the hospital and during transportation.** *Curr Opin Crit Care* 2006, **12**:43-49.
3. Norton R, Wortman E, Eastes L, Daya M, Hedges J, Hoyt J: **Appropriate helicopter transport of urban trauma patients.** *J Trauma* 1996, **41**:886-891.
4. Dunn EL, Berry PH, Cross RE Jr: **Community hospital to trauma center.** *J Trauma* 1986, **26**:733-777.
5. Reynolds HN, Habashi NM, Cottingham CA, Frawley PM, McCunn M: **Interhospital transport of the adult mechanically ventilated patient.** *Respir Care Clin N Am* 2002, **8**:37-50.
6. van Lieshout EJ, Vroom MV: **ICU Transport.** *Chest* 2005, **127**:688-689.
7. Fuchs BD, Reily DJ, Tollok E, Mallitz K, Hanson CW: **Successful Aeromedical Transport Using Inhaled Prostacyclin for a Patient With Life-Threatening Hypoxemia*.** *Chest* 2004, **125**:1579-1581.