

ORIGINAL RESEARCH

Open Access



Oxygenation strategies prior to and during prehospital emergency anaesthesia in UK HEMS practice (PREOXY survey)

Adam J. Boulton^{1,2*} , Amar Mashru³ and Richard Lyon^{3,4}

Abstract

Background: Maintaining effective oxygenation throughout the process of Pre-Hospital Emergency Anaesthesia (PHEA) is critical. There are multiple strategies available to clinicians to oxygenate patients both prior to and during PHEA. The optimal pre-oxygenation technique remains unclear, and it is unknown what techniques are being used by United Kingdom Helicopter Emergency Medical Services (HEMS). This study aimed to determine the current pre- and peri-PHEA oxygenation strategies used by UK HEMS services.

Methods: An electronic questionnaire survey was delivered to all UK HEMS services between 05 July and 26 December 2019. Questions investigated service standard operating procedures (SOPs) and individual clinician practice regarding oxygenation strategies prior to airway instrumentation (pre-oxygenation) and oxygenation strategies during airway instrumentation (apnoeic oxygenation). Service SOPs were obtained to corroborate questionnaire replies.

Results: Replies were received from all UK HEMS services ($n = 21$) and 40 individual clinicians. All services specified oxygenation strategies within their PHEA/RSI SOP and most referred to pre-oxygenation as mandatory (81%), whilst apnoeic oxygenation was mandatory in eight (38%) SOPs. The most commonly identified pre-oxygenation strategies were bag-valve-mask without PEEP (95%), non-rebreathable face mask (81%), and nasal cannula at high flow (81%). Seven (33%) services used Mapleson C circuits, whilst there were eight services (38%) that did not carry bag-valve-masks with PEEP valve nor Mapleson C circuits. All clinicians frequently used pre-oxygenation, however there was variability in clinician use of apnoeic oxygenation by nasal cannula. Nearly all clinicians (95%) reported manually ventilating patients during the apnoeic phase, with over half (58%) stating this was their routine practice. Differences in clinician pre-hospital and in-hospital practice related to availability of humidified high flow nasal oxygenation and Mapleson C circuits.

(Continued on next page)

* Correspondence: adam.boulton@nhs.net

¹Warwick Medical School, University of Warwick, Coventry, UK

²Academic Department of Anaesthesia, Critical Care, Pain and Resuscitation, Heartlands Hospital, University Hospitals Birmingham NHS Foundation Trust, Birmingham B9 5SS, UK

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

Conclusions: Pre-oxygenation is universal amongst UK HEMS services and is most frequently delivered by bag-valve-mask without PEEP or non-rebreathable face masks, whereas apnoeic oxygenation by nasal cannula is highly variable. Multiple services carry Mapleson C circuits, however many services are unable to deliver PEEP due to the equipment they carry. Clinicians are regularly manually ventilating patients during the apnoeic phase of PHEA. The identified variability in clinical practice may indicate uncertainty and further research is warranted to assess the impact of different strategies on clinical outcomes.

Keywords: Pre-hospital emergency anaesthesia, Rapid sequence induction, Pre-oxygenation, Apnoeic oxygenation, Airway management, Emergency medical services

Background

Endotracheal intubation allows definitive protection of the airway and may optimise oxygenation and ventilation in critically unwell patients [1, 2]. Advanced airway interventions, including Pre-Hospital Emergency Anaesthesia (PHEA), are being increasingly performed in the pre-hospital environment [3–6] with high procedural success [7, 8]. However, severe complications are associated with PHEA including hypoxaemia and cardiovascular instability [9–12]. One technique to minimise the risk of hypoxaemia is to provide the patient with a period of pre-oxygenation [13]. Pre-oxygenation is recommended by multiple anaesthetic guidelines in a variety of clinical settings, including pre-hospital care [14–18]. A period of successful pre-oxygenation significantly delays the onset of desaturation during apnoea by wash-out of nitrogen from the lungs, increased oxygen content in the lungs, and an increased oxygen tension in the blood and tissues [13, 19]. Patients undergoing PHEA in the pre-hospital setting are clinically unstable and regularly have airway, respiratory and haemodynamic compromise and consequently pre-oxygenation may be ineffective [20]. Given the deranged physiology, efforts to optimise pre-oxygenation are of heightened clinical significance [10, 20]. However, reliably performing adequate pre-oxygenation in critically unwell patients can be problematic due to poorly co-operative patients, concerns over gastric insufflation in an unfasted patient and equipment/logistical concerns [14, 21, 22]. There are multiple pre-oxygenation techniques available including oxygen delivery by nasal cannula, facemask, non-invasive ventilation, bag valve mask (BVM), Mapleson C circuit, or high-flow nasal cannula [11, 19, 21, 23]. The optimal technique in the pre-hospital arena is unclear and anecdotally clinicians are using a variety of techniques [14]. As such, it is uncertain which pre-oxygenation techniques are currently being employed by UK Helicopter Emergency Medical Services (HEMS) and if there is a variability between services. Although defined inconsistently in the present literature, apnoeic oxygenation may be used as an adjunct to pre-oxygenation by a range of methods [24–27]. Similarly, its application in contemporary UK HEMS practice is unknown [3]. Identification of the

oxygenation strategies being used in UK HEMS practice will allow better understanding of current practice and inform development of future observational or interventional studies. Therefore, the aim of this study was to identify the current oxygenation strategies used prior to and during PHEA by UK HEMS services and individual clinicians.

Methods

The Checklist for Reporting Results of Internet E-Surveys (CHERRIES) has been followed in the reporting of this electronic survey [28]. A cross-sectional questionnaire survey hosted by Google Forms was distributed by email to all UK HEMS services between 05 July and 26 December 2019. Questionnaires were developed to investigate service standard operating procedures (SOP) and individual clinician practice. The questionnaires were developed with pre-hospital care experts and were informed by the oxygenation strategies described in the updated Utstein-style pre-hospital advanced airway template [29]. So that responses regarding oxygenation strategies were reliable, care was taken in clearly defining oxygenation strategies through PHEA. A practical definition to limit subjectivity was used, with the point of airway instrumentation used as the point of division. Questions investigated oxygenation strategies prior to airway instrumentation (pre-oxygenation) and oxygenation strategies during airway instrumentation (apnoeic oxygenation). Piloting of the questionnaire with pre-hospital experts and iterative refinement minimised question ambiguity. The questionnaires are found in Supplementary Material 1. The first questionnaire aimed to assess the local service SOP for PHEA and oxygenation strategy described. A reply regarding the service SOP along with a copy of the service SOP to corroborate findings was sought from every UK HEMS service ($n = 21$). The second questionnaire aimed to investigate individual clinician practice. Replies were sought from two individual clinicians at each service, ideally a HEMS consultant and a HEMS junior doctor, but in services with no HEMS junior doctor, responses were sought from one senior consultant and one junior consultant.

Questions were asked in the same order to all participants and required fields ensured only complete questionnaires could be submitted.

Data were summarised using standard descriptive statistics and graphical plots. Data are reported anonymously, and individual HEMS services are not identifiable in published results. Responses were compiled and analysed using Microsoft Excel.

This survey met UK NIHR criteria as a service evaluation and therefore formal Health Research Authority approval was not required [30]. No patient identifiable data was collected.

Results

HEMS service standard operating procedures

Responses were received from every UK HEMS service ($n = 21$). A copy of the local service SOP was received from 13 services and used to corroborate questionnaire responses. A doctor-paramedic model was used by the majority of services when delivering PHEA ($n = 20$, 95%) with one service employing a paramedic-paramedic model. The approximate number of episodes of PHEA was variable across services with 12 (57%) performing > 100 per year, 6 (29%) performing 50–100 per year, and 3 (14%) < 50 per year. All services had a PHEA/RSI SOP and pre-oxygenation was referred to in this SOP by all services. No service had a separate SOP for oxygenation during PHEA. Most service SOPs referred to pre-oxygenation as mandatory ($n = 17$, 81%) and the remainder referred to pre-oxygenation as advised ($n = 4$, 19%). No SOPs identified specific patient groups in which pre-oxygenation should be used. There was a range of pre-oxygenation strategies referred to in service SOPs. The most common strategies were BVM without PEEP ($n = 20$, 95%), non-breathable face mask ($n = 17$, 81%), and nasal cannula at high flow (> 4 L/min) ($n = 17$, 81%) (Fig. 1). Specific combinations of SOP pre-oxygenation strategies varied across services (Fig. 2). The most common combination was that of BVM without PEEP, non-

rebreathable face mask, and nasal cannula at high flow (> 4 L/min) ($n = 5$, 24%). There were eight services (38%) that did not specify BVM with PEEP valve nor Mapleson C circuit within their SOP, therefore precluding delivery of PEEP. Ten service SOPs (48%) recommended use of a BVM with PEEP valve or a Mapleson C circuit, and three services (14%) identified both devices. Seven services (33%) carried Mapleson C circuits, all of whom also carried BVMs. No service SOP referred to non-invasive ventilation as a possible pre-oxygenation strategy.

Apnoeic oxygenation was discussed in all but one PHEA SOP ($n = 20$, 95%). Apnoeic oxygenation was referred to as mandatory by eight (38%), advised by seven (33%), and to be considered by 5 (24%) SOPs. Two service SOPs (10%) identified patient groups where apnoeic oxygenation should be considered, and these were obese patients and predicted airway difficulties. All service SOPs noting apnoeic oxygenation referred to nasal cannula, with 16 (76%) stating it should be at high flow (> 4 L/min) and the remainder not specifying a recommended flow.

Individual clinician practice

Doctors from every HEMS service were invited to complete questionnaires investigating their individual practice. Forty participants completed the individual questionnaire covering all HEMS services. Responses from a senior consultant and a junior doctor were received from 15 services. Four services did not have junior doctors, and hence replies were sought from one senior consultant and one junior consultant. Two services did not have a junior consultant and therefore replies were gained from one senior consultant. Questionnaire responses were received from 25 HEMS consultants and 15 HEMS junior doctors. The years of experience as a doctor ranged from 8 to 22 years (median 13, IQR 4.3) and years of experience in HEMS ranged from 1 to 20 years (median 4.5, IQR 8).

Preoxygenation strategies reported in SOPs




Fig. 1 Preoxygenation strategies reported in SOPs. BVM = Bag valve mask, PEEP = Positive End Expiratory Pressure, SOP = Standard Operating Procedure




Fig. 2 Combinations of preoxygenation strategies reported in SOPs

Thirty-six participants (90%) stated they always used pre-oxygenation and four (10%) stated they used pre-oxygenation very often (75–100% of cases). The pre-oxygenation strategies individuals reported they frequently used were similar to those identified in service SOPs (Table 1). No clinician reported use of non-invasive ventilation, in keeping with service SOPs.

Three participants (8%) stated they were not able to deliver their preferred pre-oxygenation strategy in their current service. These all related to inability to deliver PEEP and participants stated the desire to have a BVM with PEEP valve or Mapleson C available. The service SOPs these clinicians worked within showed that the service did not carry either a BVM with a PEEP valve or Mapleson C and hence PEEP could not be delivered.


Table 1 Pre-oxygenation strategies

Pre-oxygenation strategy frequently used by individuals	N	Percentage of respondents (n = 40)
Bag Valve Mask without PEEP	22	55.0%
Bag Valve Mask with PEEP	7	17.5%
Mapleson C	13	32.5%
Non-rebreathable face mask	17	42.5%
Nasal cannula, high flow (> 4 L)	17	42.5%
Nasal cannula, low flow (max 4 L)	7	17.5%
Nasal cannula, but flow not specified	1	2.5%

In contrast to pre-oxygenation, there was a large variation in how frequently individuals use apnoeic oxygenation during PHEA. There was an almost equal distribution across frequency of clinician use of apnoeic oxygenation in their last five deliveries of PHEA (Fig. 3).

Of the 35 participants (88%) who reported apnoeic oxygenation use, the majority used nasal cannula at high flow (≥ 4 L/min) (n = 32, 91%) and the remainder at low flow (max 4 L/min) (n = 3, 9%). Three participants stated that they were unable to deliver their preferred apnoeic oxygenation strategy, which was humidified high flow nasal oxygenation such as THRIVE or OptiFlow™.

We asked clinicians to describe how their oxygenation strategy changed after delivery of anaesthetic drugs but before airway instrumentation. Over half of participants (n = 23, 58%) stated they routinely ventilate patients during the apnoeic phase. A further 15 (38%) stated they would occasionally ventilate during the apnoeic phase dependent on the clinical situation, such as when the patient showed prior signs of ventilatory failure, was deemed at high risk of desaturation, or if the oxygen saturations dropped. Therefore, 95% of participants reported ventilating patients during the apnoeic phase when performing PHEA. Two participants (5%) reported no change in oxygenation strategy after delivery of anaesthetic drugs but before airway instrumentation. Twelve participants (30%) stated they increase oxygen



flow via nasal cannula from low flow to high flow (> 4 L/min).

When comparing their overall oxygenation strategies between PHEA and in-hospital practice, 14 participants (35%) reported there was no difference. In the remainder, differences between PHEA and in-hospital technique were use of humidified high flow nasal oxygenation such as THRIVE or OptiFlow ($n = 14$, 35%), and use of a Mapleson C circuit ($n = 14$, 35%).

Discussion

Our study set out to provide a snapshot of current PHEA practices across UK HEMS services, with a particular focus on pre-oxygenation and apnoeic oxygenation strategies. When evaluating the merits, drawbacks and lessons learned from this study, it is worth noting initially that the survey was answered by every HEMS in the UK. This provides us not only with a valuable insight into the makeup of our HEMS around the country, their utilisation of SOPs and their rates of PHEA, but as a springboard for sharing, comparing and understanding practice such that it can be enhanced for the benefit of the specialty and the patients it serves.

More than 100 PHEA episodes were undertaken each year by the majority of services, with PHEA being detailed by every service in a dedicated SOP. Pre-oxygenation strategies were widely implemented during PHEA and were mentioned in every SOP. However, it is worth noting that 20% of services did not mandate its use. Individuals appeared more universally keen to

utilise pre-oxygenation, but it was still not implemented 100% of the time in those surveyed. This study is not intended to debate the pros and cons of such practice, instead we see this as a valuable example of the purpose of work such as ours in challenging or supporting assumptions that may exist regarding PHEA. We noted a spread of modalities for delivering preoxygenation are currently utilised, with very few respondents feeling they were unable to deliver their preferred pre-oxygenation strategy with the pre-hospital equipment, or indeed the SOP they were working with. However, where discordance did exist it seemed to regard the utilisation of PEEP, either through BVM or Mapleson C. With many services providing the option of PEEP and around a third suggesting availability of a Mapleson C circuit, it raises stimulating discussion regarding variation in equipment availability, governance and oxygenation strategies amongst different services and whether these should exist. The carriage of Mapleson C circuits could be related to clinician familiarity and tactile feedback in terms of delivering PEEP and manual ventilation. It should be noted that all services carrying Mapleson C circuits also carried BVMs.

Although apnoeic oxygenation strategies were mentioned in all but one service SOP, recommendations for apnoeic oxygenation strategies appeared far less firm than for pre-oxygenation. In contrast to pre-oxygenation, a majority of services did not mandate the use of apnoeic oxygenation during PHEA. This may reflect the current evidence base for apnoeic oxygenation and its practical difficulties in the pre-hospital environment [24, 31]. It is interesting to note that many strategies advised nasal cannula with flow rates > 4 L/min and many clinicians reported their preference desire to deliver humidified high flow nasal oxygen, whilst no clinician or SOP reported use of non-invasive ventilation. All these strategies have resource implications, most significantly high flow nasal oxygen and non-invasive ventilation, requiring large amounts of oxygen and equipment that may limit their application to the pre-hospital environment. This heterogeneity in clinical practice highlights an area of uncertainty that merits further research and discussion.

Notably, 95% of clinicians reported manually ventilating patients during the apnoeic phase of PHEA. This practice is at odds with the description of traditional RSI [32] and the practice of many in-hospital clinicians [33]. Manual ventilation during RSI has remained a contentious topic, with some believing avoidance of mask ventilation is dogmatic practice [34]. There is mounting evidence in favour of mask ventilation in RSI [35], and this is reflected in practice guidelines [16]. The current UK pre-hospital practice revealed by this survey may be due to rapid

translation of research into practice and clinician desire to practice evidence-based medicine.

This desire to capture and understand the reality of UK practice was reflected in our study design and mirrored in the pragmatic approach that we took to pre-oxygenation and apnoeic oxygenation definitions. Present literature definitions of apnoeic oxygenation are variable and imprecise [14, 16, 24, 36]. Often, apnoeic oxygenation is defined as oxygenation post administration of muscle relaxant during RSI [16, 36]. However, real-world application involves a secondary supply of oxygen, through an alternative route, usually by nasal cannula that remains in place during airway instrumentation [24, 31]. For this reason, our questions were targeted to oxygenation strategies pre and per airway instrumentation. Our findings suggest that many practitioners and services had a preference for two modalities of oxygen delivery pre-airway instrumentation, with one modality remaining in place during airway instrumentation to deliver apnoeic oxygenation. Better understanding of practical delivery of PHEA oxygenation and clarification of this nuance may assist in future description, comparison of practice, and standardisation of future nomenclature.

Our study achieved an excellent response rate covering all UK HEMS services. The results therefore likely represent a good summary of contemporaneous practice. The survey methodology presents risks of social acceptability and recall bias, however obtaining service SOP copies where possible helped to mitigate this. Nevertheless, participant responses may not represent real-life clinical practice. Whilst this survey has provided a comprehensive assessment of current UK practice, it does not assess clinical outcomes and hence further clinical studies investigating patient outcomes are needed. The heterogeneity in clinical practice highlighted by this study emphasises that the optimal pre-oxygenation strategy remains unclear and there appears to be uncertainty in the value of apnoeic oxygenation, and hence more detailed clinical studies are warranted.

Without understanding existing clinical practice we are unable to identify where practice is routine or divergent, and therefore where to prioritise research and focus to improve patient care. Whilst it is clearly important that individual services need to tailor care to meet their patients' needs and, within those services, clinicians may need the flexibility to tailor strategies to meet the needs of each individual patient, we believe our study contributes to the national dialogue, highlights the benefits of sharing data and practice and thus, ultimately, allows shared learning across the speciality.

Conclusions

This survey of all UK HEMS services identifies that pre-oxygenation is almost universal in current PHEA

practice and most frequently delivered by bag valve mask without PEEP or non-rebreathable face masks. Conversely, use of apnoeic oxygenation is not widespread and there is significant variability in both individual clinician practice and service SOPs. Many clinicians are routinely ventilating patients during the apnoeic phase, whilst others are ventilating only in high risk patients. Differences between in-hospital and pre-hospital practice were principally due to availability of equipment such as humidified high flow nasal oxygenation and Mapleson C circuits. Multiple services carry bag valve masks with PEEP and/or Mapleson C circuits, however many services carry neither therefore precluding delivery of PEEP. All those carrying Mapleson C circuits also carried bag valve masks with or without a PEEP valve. The identified variability in oxygenation strategies during PHEA warrants further study, including assessment of clinical endpoints.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13049-020-00794-x>.

Additional file 1.

Abbreviations

BVM: Bag Valve Mask; HEMS: Helicopter Emergency Medical Service; PEEP: Positive End Expiratory Pressure; PHEA: Prehospital Emergency Anaesthesia; RSI: Rapid Sequence Induction; SOP: Standard Operating Procedure; THRIVE: Transnasal Humidified Rapid-Insufflation Ventilatory Exchange

Acknowledgments

We sincerely thank the participants for taking the time to diligently complete the questionnaires.

Authors' contributions

Study design: AJB, AM, RL. Study conduct: AJB, AM. Data analysis and figure production: AJB. Drafting of manuscript: AJB, AM. Review of manuscript: AJB, AM, RL. The authors read and approved the final manuscript.

Funding

No specific funding was obtained to support this study.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

This survey met UK NIHR criteria as a service evaluation and therefore formal Health Research Authority approval was not required [30]. No patient identifiable data was collected.

Consent for publication

All authors provide consent for publication.

Competing interests

Authors identify no conflicts of interest.

Author details

¹Warwick Medical School, University of Warwick, Coventry, UK. ²Academic Department of Anaesthesia, Critical Care, Pain and Resuscitation, Heartlands Hospital, University Hospitals Birmingham NHS Foundation Trust, Birmingham B9 5SS, UK. ³Air Ambulance Kent Surrey Sussex, Rochester

Airport, Maidstone Road, Chatham, Rochester, UK. ⁴University of Surrey, Stag Hill, Guildford, UK.

Received: 12 August 2020 Accepted: 7 October 2020

Published online: 12 October 2020

References

- Lockey DJ, Healey B, Crewdson K, Chalk G, Weaver AE, Davies GE. Advanced airway management is necessary in prehospital trauma patients. *Br J Anaesth*. 2015;114(4):657–62.
- Sise MJ, Shackford SR, Sise CB, Sack DI, Paci GM, Yale RS, et al. Early intubation in the management of trauma patients: indications and outcomes in 1,000 consecutive patients. *J Trauma*. 2009;66(1):32–9 discussion 9–40.
- Crewdson K, Rehn M, Lockey D. Airway management in pre-hospital critical care: a review of the evidence for a 'top five' research priority. *Scand J Trauma Resusc Emerg Med*. 2018;26(1):89.
- Hubble MW, Brown L, Wilfong DA, Hertelendy A, Benner RW, Richards ME. A meta-analysis of prehospital airway control techniques part I: orotracheal and nasotracheal intubation success rates. *Prehosp Emerg Care*. 2010;14(3):377–401.
- Crewdson K, Lockey DJ, Roislien J, Lossius HM, Rehn M. The success of pre-hospital tracheal intubation by different pre-hospital providers: a systematic literature review and meta-analysis. *Critical Care (London, England)*. 2017; 21(1):31.
- Sunde GA, Sandberg M, Lyon R, Fredriksen K, Burns B, Hufthammer KO, et al. Hypoxia and hypotension in patients intubated by physician staffed helicopter emergency medical services - a prospective observational multi-Centre study. *BMC Emerg Med*. 2017;17(1):22.
- Lockey DJ, Crewdson K. Pre-hospital anaesthesia: no longer the 'poor relative' of high quality in-hospital emergency airway management. *Br J Anaesth*. 2018;120(5):898–901.
- Gellerfors M, Fevang E, Backman A, Kruger A, Mikkelsen S, Nurmi J, et al. Pre-hospital advanced airway management by anaesthetist and nurse anaesthetist critical care teams: a prospective observational study of 2028 pre-hospital tracheal intubations. *Br J Anaesth*. 2018;120(5):1103–9.
- Lockey D, Crewdson K, Weaver A, Davies G. Observational study of the success rates of intubation and failed intubation airway rescue techniques in 7256 attempted intubations of trauma patients by pre-hospital physicians. *Br J Anaesth*. 2014;113(2):220–5.
- Schwartz DE, Matthay MA, Cohen NH. Death and other complications of emergency airway management in critically ill adults. A prospective investigation of 297 tracheal intubations. *Anesthesiology*. 1995;82(2):367–76.
- Davis DP, Hwang JQ, Dunford JV. Rate of decline in oxygen saturation at various pulse oximetry values with prehospital rapid sequence intubation. *Prehosp Emerg Care*. 2008;12(1):46–51.
- Sunde GA, Heltne JK, Lockey D, Burns B, Sandberg M, Fredriksen K, et al. Airway management by physician-staffed Helicopter Emergency Medical Services - a prospective, multicentre, observational study of 2,327 patients. *Scand J Trauma Resusc Emerg Med*. 2015;23:57.
- Nimmagadda U, Salem MR, Crystal GJ. Preoxygenation: physiologic basis, benefits, and potential risks. *Anesth Analg*. 2017;124(2):507–17.
- Lockey DJ, Crewdson K, Davies G, Jenkins B, Klein J, Laird C, et al. AAGBI: safer pre-hospital anaesthesia 2017: Association of Anaesthetists of Great Britain and Ireland. *Anaesthesia*. 2017;72(3):379–90.
- Frerk C, Mitchell VS, McNarry AF, Mendonca C, Bhagrath R, Patel A, et al. Difficult airway society 2015 guidelines for management of unanticipated difficult intubation in adults. *Br J Anaesth*. 2015;115(6):827–48.
- Higgs A, McGrath BA, Goddard C, Rangasami J, Suntharalingam G, Gale R, et al. Guidelines for the management of tracheal intubation in critically ill adults. *Br J Anaesth*. 2018;120(2):323–52.
- Mushambi MC, Kinsella SM, Popat M, Swales H, Ramaswamy KK, Winton AL, et al. Obstetric Anaesthetists' Association and difficult airway society guidelines for the management of difficult and failed tracheal intubation in obstetrics. *Anaesthesia*. 2015;70(11):1286–306.
- Crewdson K, Lockey D, Voelckel W, Temesvari P, Lossius HM, Group EMW. Best practice advice on pre-hospital emergency anaesthesia & advanced airway management. *Scand J Trauma Resusc Emerg Med*. 2019;27(1):6.
- Weingart SD, Levitan RM. Preoxygenation and prevention of desaturation during emergency airway management. *Ann Emerg Med*. 2012;59(3):165–75 e1.
- Mort TC. Preoxygenation in critically ill patients requiring emergency tracheal intubation. *Crit Care Med*. 2005;33(11):2672–5.
- Weingart SD, Trueger NS, Wong N, Scofi J, Singh N, Rudolph SS. Delayed sequence intubation: a prospective observational study. *Ann Emerg Med*. 2015;65(4):349–55.
- Lawes EG, Campbell I, Mercer D. Inflation pressure, gastric insufflation and rapid sequence induction. *Br J Anaesth*. 1987;59(3):315–8.
- Frat JP, Ricard JD, Coudroy R, Robert R, Ragot S, Thille AW. Preoxygenation with non-invasive ventilation versus high-flow nasal cannula oxygen therapy for intubation of patients with acute hypoxaemic respiratory failure in ICU: the prospective randomised controlled FLORALI-2 study protocol. *BMJ Open*. 2017;7(12):e018611.
- Wimalasena Y, Burns B, Reid C, Ware S, Habig K. Apneic oxygenation was associated with decreased desaturation rates during rapid sequence intubation by an Australian helicopter emergency medicine service. *Ann Emerg Med*. 2015;65(4):371–6.
- Eross A, Hetzman L, Petroczy A, Gorove L. Apneic preoxygenation without nasal prongs: the "Hungarian Air Ambulance method". *Scand J Trauma Resusc Emerg Med*. 2016;24(1):5.
- Ramachandran SK, Cosnowski A, Shanks A, Turner CR. Apneic oxygenation during prolonged laryngoscopy in obese patients: a randomized, controlled trial of nasal oxygen administration. *J Clin Anesth*. 2010;22(3):164–8.
- Patel A, Nouraei SAR. Transnasal humidified rapid-insufflation Ventilatory exchange (THRIVE): a physiological method of increasing apnoea time in patients with difficult airways. *Anaesthesia*. 2015;70(3):323–9.
- Eysenbach G. Improving the quality of web surveys: the checklist for reporting results of internet E-surveys (CHERRIES). *J Med Internet Res*. 2004; 6(3):e34.
- Sunde GA, Kottmann A, Heltne JK, Sandberg M, Gellerfors M, Kruger A, et al. Standardised data reporting from pre-hospital advanced airway management - a nominal group technique update of the Utstein-style airway template. *Scand J Trauma Resusc Emerg Med*. 2018;26(1):46.
- Authority HR. Defining Research Table 2017 [Available from: http://www.hra-decisiontools.org.uk/research/docs/DefiningResearchTable_Oct2017-1.pdf].
- Semler MW, Janz DR, Lentz RJ, Matthews DT, Norman BC, Assad TR, et al. Randomized trial of apneic oxygenation during endotracheal intubation of the critically ill. *Am J Respir Crit Care Med*. 2016;193(3):273–80.
- Stept WJ, Safar P. Rapid induction-intubation for prevention of gastric-content aspiration. *Anesth Analg*. 1970;49(4):633–6.
- Sajayan A, Wicker J, Ungureanu N, Mendonca C, Kimani PK. Current practice of rapid sequence induction of anaesthesia in the UK - a national survey. *Br J Anaesth*. 2016;117(suppl_1):i69–74.
- Kritek PA, Luks AM. Preventing dogma from driving practice. *N Engl J Med*. 2019;380(9):870–1.
- Casey JD, Janz DR, Russell DW, Vonderhaar DJ, Joffe AM, Dischert KM, et al. Bag-mask ventilation during tracheal intubation of critically ill adults. *N Engl J Med*. 2019;380(9):811–21.
- Patel A, Nouraei SA. Transnasal humidified rapid-insufflation Ventilatory exchange (THRIVE): a physiological method of increasing apnoea time in patients with difficult airways. *Anaesthesia*. 2015;70(3):323–9.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

